



Randomness Zoo

Antoine Taveneaux

► To cite this version:

| Antoine Taveneaux. Randomness Zoo. 2012. hal-00850992

HAL Id: hal-00850992

<https://hal.science/hal-00850992>


Preprint submitted on 10 Aug 2013

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Randomness Zoo

Antoine Tavenaux



Definition 1 (MLR: Martin-Löf Randomness (Definition 3.2.1 in [ML66])).

1. A Martin-Löf test, a ML-test for short, is a uniformly c.e. sequence $(G_m)_{m \in \mathbb{N}}$ of open sets such that $\forall n \mu(G_m) \leq 2^{-m}$.
2. A set R fails the test if $R \in \cap_{m \in \mathbb{N}} G_m$ otherwise R passes the test.
3. R is Martin-Löf Random if R passes each ML-test.

1 Weaker than MLR

Definition 2 (Scan rule function). For a partial function $f : 2^{<\omega} \rightarrow \{\text{scan}, \text{select}\} \times \mathbb{N}$ we denote $n : 2^{<\omega} \rightarrow \mathbb{N}$ and $\delta : 2^{<\omega} \rightarrow \{\text{scan}, \text{select}\}$.

And we say that f is a scan rule if for all $\sigma, \rho \in 2^{<\omega}$ such that $\sigma \prec \rho$ we have $n(\sigma) \neq n(\rho)$.

The sequence of string observed by f on $Z \in cs$ is defined by (if f is well defined on this points):

$$\begin{aligned} V_f^A(0) &= A(n(\varepsilon)) \\ V_f^A(k+1) &= V_f^A(k).A(n(V_f^A(k))) \end{aligned}$$

and bits selected by f are:

$$\begin{aligned} T_f^A(0) &= \varepsilon \\ T_f^A(n+1) &= \begin{cases} T_f^A(n) & \text{if } \delta(V_f^A(n)) = \text{scan} \\ T_f^A(n).V_f^A(n+1) & \text{if } \delta(V_f^A(n)) = \text{select} \end{cases} \end{aligned}$$

and we say that f is well defined on Z if $V_f^A(k)$ is well define for all k and $(T_f^A(n))$ converge to an infinite string and we denote this infinite string by T_f^A .

Definition 3 (Martingale). A martingale is a function $\mathcal{M} : 2^{<\omega} \rightarrow \mathbb{R}^+ \cup 0$ such that for all $\sigma \in 2^{<\omega}$

$$\mathcal{M}(\sigma) = \frac{\mathcal{M}(\sigma 0) + \mathcal{M}(\sigma 1)}{2}$$

We say that \mathcal{M} succeed on $Z \in 2^\omega$ if

$$\limsup_{n \rightarrow \infty} \mathcal{M}(Z \upharpoonright n) = \infty$$

Definition 4 (KLR: KolmogorovLoveland randomness (Definition in [Kol63] and [Lov66])). $R \in 2^\omega$ is KL-random if for any partial computable scan rule function f and any partial computable martingale \mathcal{M} such that f is well defined on R the martingale \mathcal{M} does not succeed on T_f^A .

Definition 5 (KLStoch: KolmogorovLoveland stochasticity (Definition in [Lov66])). Let $\#0 : 2^{<\omega} \rightarrow \mathbb{N}$ the function giving the number of “0” in a string.

A sequence R is KolmogorovLoveland stochastic if for all partial computable scan rule f such that f is well defined on Z we have:

$$\lim_{n \rightarrow \infty} \frac{\#0(T_f^A \upharpoonright n)}{n} = \frac{1}{2}$$

Definition 6 (MWCStoch: Mises-Wald-Church stochasticity (Definition in [vM19] and [Chu40])). A sequence R is Mises-Wald-Church stochastic if for all partial computable monotonic scan rule function f is well defined on Z we have:

$$\lim_{n \rightarrow \infty} \frac{\#0(T_f^A \upharpoonright n)}{n} = \frac{1}{2}$$

Definition 7 (ChStoch: Church stochasticity (Definition in [vM19] and [Chu40])). A sequence R is Church stochastic if for all total computable monotonic scan rule function f we have:

$$\lim_{n \rightarrow \infty} \frac{\#0(R_{f(0)}R_{f(1)} \dots R_{f(n)})}{n} = \frac{1}{2}$$

Definition 8 (PIInjR: partial injective randomness (Definition in [MN06])).

A sequence R is partial injective random if for any total computable injective function $g : \mathbb{N} \rightarrow \mathbb{N}$ and any partial computable martingale \mathcal{M} this martingale is defined and does not succeed on the sequence $R_{f(1)}R_{f(2)} \dots R_{f(n)}R_{f(n+1)} \dots$.

Definition 9 (InjR: injective randomness (Definition in [?])).

A sequence R is injective random if for any total computable injective function $g : \mathbb{N} \rightarrow \mathbb{N}$ and any total computable martingale \mathcal{M} this martingale does not succeed on the sequence $R_{f(1)}R_{f(2)} \dots R_{f(n)}R_{f(n+1)} \dots$.

Definition 10 (PermR: partial permutation randomness (Definition in [BHKM09])).

A sequence R is partial permutation random if for any total computable bijective function $g : \mathbb{N} \rightarrow \mathbb{N}$ and any partial computable martingale \mathcal{M} this martingale is defined and does not succeed on the sequence $R_{f(1)}R_{f(2)} \dots R_{f(n)}R_{f(n+1)} \dots$.

Definition 11 (PCR: partial computable randomness (Definition in [AS97])).

A sequence R is partial computable random if for all partial computable martingale \mathcal{M} if $\mathcal{M}(R \upharpoonright n)$ is define for all n and \mathcal{M} does not succeed on R .

Definition 12 (CR: computable randomness (Definition in [Sch71])). A sequence R is computable random if for all total computable martingale \mathcal{M} this martingale succeed on R .

Definition 13 (SR: Schnorr randomness (Definition in [Sch71])). A Schnorr test is a uniformly c.e. sequence $(G_m)_{m \in \mathbb{N}}$ of open sets such that $\forall n \mu(G_m) = 2^{-m}$.

R is Schnorr random if for any Schnorr test $(G_m)_{m \in \mathbb{N}}$ $R \notin \bigcap_{m \in \mathbb{N}} G_m$

Definition 14 (FBoundR: finitely bounded randomness (Definition in [BDN12])). R is finitely bounded random if R passes any Martin-Löf test (U_n) such that for every n , $\#U_n < \infty$ (with $\#U_n$ the number of sting enumerated in U_n).

Definition 15 (CBoundR: computably bounded randomness (Definition in [BDN12])). A Martin-Löf test (U_n) is computably bounded if there is some total computable function f such that $\#U_n \leq f(n)$ for every n .

R is computably bounded random if R passes every computably bounded Martin-Löf test.

Definition 16 (WR: weakly randomness (Definition in [Kur81])). R is weakly random if $R \in U$ for every Σ_1^0 set $U \subseteq 2^\omega$ of measure 1.

Definition 17 (PolyR: polynomial randomness (Definition in [Wan96])). A sequence R is polynomially random if any martingale computable in polynomial time \mathcal{M} do not succeed on R .

Definition 18 ($\text{dim}_{\text{comp}}^s \mathbf{R}$: computable s -randomness (Definition in [Lut03] and [May02])). A computable s -test is a uniformly computable sequence $(G_m)_{m \in \mathbb{N}}$ of computable open sets such that for all n

$$\sum_{x \in G_m} 2^{-s|x|} \leq 2^{-n}.$$

R is computably s -random if for all $s' < s$ and computable s -tests (G_m) we have:

$$R \notin \bigcap_{m \in \mathbb{N}} G_m$$

Definition 19 ($\text{Cdim}^s \mathbf{R}$: constructive s -randomness (Definition in [Lut03] and [May02])). A constructive s -test is a uniformly computable sequence $(G_m)_{m \in \mathbb{N}}$ of computable enumerable open sets such that for all n

$$\sum_{x \in G_m} 2^{-s|x|} \leq 2^{-n}.$$

R is computably s -random if for all $s' < s$ and computable s -tests (G_m) we have:

$$R \notin \bigcap_{m \in \mathbb{N}} G_m$$

2 Stronger than MLR

Definition 20 (DiffR: difference randomness (Definition in [FN])). A difference test is given by a sequence $(V_m)_{m \in \mathbb{N}}$ of uniformly c.e. sets and a Π_1^0 set P such that $\mu(P \cap V_m) \leq 2^{-m}$ for every m .

A sequence R is difference random if for any difference test $(V_m)_{m \in \mathbb{N}}, P$ we have

$$R \notin P \cap \left(\bigcap_{m \in \mathbb{N}} V_m \right).$$

Definition 21 (BalancedR: balanced randomness (Definition in [FHM⁺10])).

A balanced test is a sequence $(V_m)_{m \in \mathbb{N}}$ of c.e. sets such that $V_i = W_{f(i)}$ for some 2^n -c.e. function f and $\mu(V_m) \leq 2^{-m}$ for every m .

A sequence R is balanced random if R passes any balanced test.

Definition 22 (WDemR: weak Demuth randomness (Definition in [Dem82])).

Demuth test is a sequence $(V_m)_{m \in \mathbb{N}}$ of c.e. sets such that $V_i = W_{f(i)}$ for some ω -c.e. function f and $\mu(V_m) \leq 2^{-m}$ for every m .

A sequence R is weak Demuth random if for any Demuth test (V_m) we have $R \notin \bigcap_{m \in \mathbb{N}} V_m$.

Definition 23 (DemR: Demuth randomness (Definition in [Dem82])). A

Demuth test is a sequence $(V_m)_{m \in \mathbb{N}}$ of c.e. sets such that $V_i = W_{f(i)}$ for some ω -c.e. function f and $\mu(V_m) \leq 2^{-m}$ for every i .

A sequence R is Demuth random if for any Demuth test (V_m) we have $R \notin V_m$ for almost all m .

Definition 24 (W2R: weak 2-randomness (Definition in [Kau91])). A sequence R is weak 2-random if $R \notin U$ for every Π_2^0 set $U \subset 2^\omega$ of measure 0.

Definition 25 (LimitR: limit randomness (Definition in [KN11])). A limit test is a sequence $(V_m)_{m \in \mathbb{N}}$ of c.e. sets such that $V_i = W_{f(i)}$ for some Δ_2^0 -computable function f and $\mu(V_m) \leq 2^{-m}$ for every m .

A sequence R is limit random if for any limit test (V_m) we have $R \notin V_m$ for almost all m .

Definition 26 (Δ_1^1 R: Δ_1^1 randomness (Definition in [ML70])). R is Δ_1^1 -random if R avoids each null Δ_1^1 -class.

Definition 27 ($\Pi_1^1\mathbf{MLR}$: Π_1^1 -Martin-Löf Randomness (Definition in [HN07])).
A Π_1^1 -Martin-Löf test is a sequence $(G_m)_{m \in \mathbb{N}}$ of open sets such that $\forall n \ \mu(G_m) \leq 2^{-m}$ and the relation $\{\langle m, \sigma \rangle \mid [\sigma] \subseteq G_m\}$ is Π_1^1 .
 R is Π_1^1 -Martin-Löf Random if R passes each Π_1^1 -ML-test.

Definition 28 ($\Pi_1^1\mathbf{R}$: Π_1^1 -Randomness (Definition in [HN07])). R is Π_1^1 -random if R avoids each null Π_1^1 -class.

References

- [AS97] K. Ambos-Spies. Algorithmic randomness revisited. In *Language, Logic and Formalization of Knowledge*, pages 33–52, 1997.
- [BDN12] P. Brodhead, R. Downey, and K. M. Ng. Bounded randomness. In *Computation, Physics and Beyond*, pages 59–70, 2012.
- [BHKM09] L. Bienvenu, R. Hölzl, T. Kräling, and W. Merkle. Separations of non-monotonic randomness notions. In *6th Int’l Conf. on Computability and Complexity in Analysis*, 2009.
- [Chu40] A. Church. On the concept of a random sequence. *Bull. Amer. Math. Soc.*, 46:130–135, 1940.
- [Dem82] O. Demuth. Some classes of arithmetical real numbers. *Comment. Math. Univ. Carolin.*, 23(3):453–465, 1982.
- [FHM⁺10] S. Figueira, D. Hirschfeldt, J. Miller, Selwyn Ng, and A Nies. Counting the changes of random Δ_2^0 sets. In *CiE 2010*, pages 1–10, 2010. Journal version to appear in J.Logic. Computation.
- [FN] J. Franklin and K. M. Ng. Difference randomness. *Proceedings of the American Mathematical Society*. To appear.
- [HN07] G Hjorth and A Nies. Randomness in effective descriptive set theory. *J. London. Math. Soc.*, 75(2):495–508, 2007.
- [Kau91] S. Kautz. *Degrees of random sets*. Ph.D. Dissertation, Cornell University, 1991.
- [KN11] A. Kučera and A Nies. Demuth randomness and computational complexity. *Ann. Pure Appl. Logic*, 162:504–513, 2011.
- [Kol63] A. N. Kolmogorov. On tables of random numbers. *Sankhyā Ser. A*, 25:369–376, 1963. Reprinted in Theoret. Comput. Sci. 207(2) (1998) 387-395.
- [Kur81] S. Kurtz. *Randomness and genericity in the degrees of unsolvability*. Ph.D. Dissertation, University of Illinois, Urbana, 1981.
- [Lov66] D. Loveland. A new interpretation of the von Mises’ concept of random sequence. *Z. Math. Logik Grundlagen Math.*, 12:279–294, 1966.

- [Lut03] J. H. Lutz. The dimensions of individual strings and sequences. *Information and Computation*, 187, 2003.
- [May02] Elvira Mayordomo. A Kolmogorov complexity characterization of constructive Hausdorff dimension. *Inform. Process. Lett.*, 84(1):1–3, 2002.
- [ML66] P. Martin-Löf. The definition of random sequences. *Inform. and Control*, 9:602–619, 1966.
- [ML70] P. Martin-Löf. On the notion of randomness. In *Intuitionism and Proof Theory (Proc. Conf., Buffalo, N.Y., 1968)*, pages 73–78. North-Holland, Amsterdam, 1970.
- [MN06] J. Miller and A. Nies. Randomness and computability: Open questions. *Bull. Symbolic Logic*, 12(3):390–410, 2006.
- [Sch71] C.P. Schnorr. *Zufälligkeit und Wahrscheinlichkeit. Eine algorithmische Begründung der Wahrscheinlichkeitstheorie*. Springer-Verlag, Berlin, 1971. Lecture Notes in Mathematics, Vol. 218.
- [vM19] R. von Mises. Grundlagen der Wahrscheinlichkeitsrechnung. *Math. Zeitschrift*, 5:52–99, 1919.
- [Wan96] Yongge Wang. *Randomness and Complexity*. PhD dissertation, University of Heidelberg, 1996.